

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS**  
**FOR ANIMAL TREATMENT ONLY**

**PASTORAL / AG**™

# IMIDALAB™

## POUR-ON

### LICE TREATMENT FOR SHEEP

**ACTIVE CONSTITUENT: 35 g/L IMIDACLOPRID**  
**Also contains: 300 g/L N-METHYL-2-PYRROLIDONE (solvent)**

For the control of neonicotinoid susceptible body lice *Bovicola ovis* on shorn sheep up to 7 days off-shears and unshorn lambs up to 2 months of age. This product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hrs of shearing.

Resistance can develop to any chemical.



**20 L**

## DIRECTIONS FOR USE



### Restraints:

**DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.**

### Dosage and Administration:

This product can be used on all adult sheep up to 7 days off-shears and on unshorn lambs up to 2 months of age.

Apply the entire dose to sheep under 30 kg as a single broadband stripe from poll to tail butt. For sheep over 30 kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the butt of the tail. The stripes should be no further than 10 cm apart. Refer to the diagrams below.

Bodyweight (kg)	Dose (mL)	Method	Apply in the pattern below
Below 6 kg	8	Single Stripe	
6 - 8	12		
8.1 - 10	15		
10.1 - 12.5	20		
12.6 - 15	25		
15.1 - 20	30		
20.1 - 30	40	Double Stripe	
30.1 - 55	60		
55.1 - 80	80		
80.1 - 100	90		

Add 10 mL for each 10 kg above 100 kg.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group (e.g. lambs), dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

### General Directions:

Apply product only with a 6-stripe applicator.

To prevent reinfestation treated sheep must not be mixed with lice-free sheep until 6 weeks after treatment. Live lice on ewes treated within 6 weeks of lambing can infect lambs from these ewes. Treated sheep must not be mixed with lousy sheep. Do not use on sheep which have not been cleanly shorn for any reason e.g. mycotic dermatitis.

Sheep must be treated within 7 days of shearing. Unshorn lambs can be treated up to 2 months of age. Efficacy is not affected by moderate rainfall either before or after treatment (50 mm rain 1 hour prior to treatment and/or 12.5 mm 2 hours after treatment and/or 25 mm 24 hours after treatment).

Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control.

Treated sheep could potentially develop backline

dermatitis following prolonged exposure to extreme heat and sunlight without access to shade. Do not leave container, hose or applicator in the sun for long periods of time.

PASTORAL AG Imidalab Pour-On may be used in conjunction with worm control drenches and at the same time as vaccination.

PASTORAL AG Imidalab Pour-On contains a blue dye to identify treated sheep. This dye is scourable from the wool. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces.

PASTORAL AG Imidalab Pour-On is safe for use on breeding rams.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 21 days before slaughter for human consumption.**

**WOOL: DO NOT USE less than 2 months before shearing or fibre collection.**

### TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI)

**DO NOT USE less than 63 days before slaughter for export. Before using this product, confirm the current ESI from the distributor on (03) 5820 8444 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**

### SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container, wear face shield. Wash hands after use. After each day's use, wash face shield.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. If in eyes wash out immediately with water.

### ADDITIONAL USER SAFETY INFORMATION

Rehandling Interval: Do not rehandle sheep after treatment until dry.

Additional information is listed in the safety data sheet.

### ENVIRONMENTAL PROTECTION

Imidacloprid is highly toxic to aquatic species. Do not contaminate dams, ponds, rivers, streams or other waterways with the chemical or used container.

### DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be recycled separately with the container. If not recycling, break, crush or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with local, state or territory government regulations. Do not burn empty containers or product.

### STORAGE

Store below 30°C (room temperature) and protect from light in a closed, original container.

**Causes skin and eye irritation.  
May cause respiratory irritation.  
May damage the unborn child.**

Avoid breathing fumes, mists, vapours or spray.

Do not get in eyes, on skin, or on clothing.

Use only outdoors or in a well ventilated area.

Wear protective gloves, protective clothing and eye or face protection.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

B:

EXP:



26710B-7