

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**

PASTORAL / AG™

STRIKELAB®

SPRAY-ON SHEEP BLOWFLY TREATMENT

ACTIVE CONSTITUENT: 50 g/L DICYCLANIL

For the protection of sheep, either off-shears or with any length wool, against strike by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*) for 18 to 24 weeks.

For the protection of mulesing and marking wounds on sheep against strike by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*) during the wound healing process.

PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment contains dicyclanil, a member of the pyrimidine class of chemicals. It is effective against sensitive strains of blowfly (*Lucilia cuprina*). Resistance may develop to any chemical.



20 L

DIRECTIONS FOR USE:**Restraints:**

DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.
DO NOT re-treat animals for a minimum of 63 days after last treatment.

Dosage and Administration:**SHAKE WELL BEFORE USE.**

PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment is applied as a low-volume treatment on to areas where protection is required. When applying PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment, the nozzle of the applicator gun should be high enough above the fleece (approximately 15 - 25 cm), to achieve a 15 cm wide band at each pass for protection of body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool.

DO NOT apply during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for mulesing/marking to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis.

FOR THE PREVENTION OF BODY STRIKE (Diagram 1)			
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
10-20	14	7	2
21-30	16	8	2
31-50	20	10	2
>50	24	12	2

FOR THE PREVENTION OF BREECH STRIKE (Diagram 1)			
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
10-20	7	7	1
21-30	8	8	1
31-50	10	10	1
>50	12	12	1

FOR THE PREVENTION OF POLL STRIKE (Diagram 1)
 Rams and sheep susceptible to poll strike should be treated. Apply 5 mL of PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment to the areas requiring protection, around the poll and horns.

FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (Diagram 2)
 Apply two bands, one over each side of the mulesing wound and overlapping 25 mm onto the surrounding wool.


SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
Up to 30	8	4	2
31-50	10	5	2

FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (Diagram 3)
 When a tail has been amputated, apply 4 mL over the crutch ensuring sufficient coverage of the open wound.
 When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied.
 In the case of ram lambs, a further 4 mL can be sprayed over and around the scrotum area after the rubber ring has been applied.

LAMBS WEIGHT (> 30 kg)	TOTAL VOLUME / LAMB (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS
Ram lambs	8	4	2
Ewe lambs	4	4	1

CRITICAL COMMENTS

1. Use only a spray-on applicator gun recommended by the manufacturer.
2. Ensure the applicator gun is clean.
3. Calibrate applicator gun using an accurate volume measure.
4. Ensure all areas to be protected are treated.
5. When preventing body strike ensure bands overlap along the midline – see Diagram 1.
6. When protecting mulesing wound from fly strike, take special care to treat the wound and the surrounding 25 mm of wool – see Diagram 2.
7. When protecting marking wounds from fly strike, take special care to treat the wound and surrounding wool – see Diagram 3.
8. This product is not recommended for treating existing strikes. Use a product registered as a dressing for existing strikes.
9. Heavy rain following application of this product could diminish the period of protection.
10. PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment contains a scorable colouring agent for easy band identification.
11. Withholding periods must be observed following application of PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment for any purpose including after mulesing / marking as finite residues will occur.

**General Directions:**

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment provides season long protection to sheep against body, breach and poll strike and may be used either off-shears or on sheep with any length wool. PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment also protects mulesing and marking wounds on lambs against fly strike during the wound healing process. The dose volume is determined by the weight of sheep being treated. Apply PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment according to the diagrams and tables in the Directions For Use.

DO NOT dilute. DO NOT mix with any other product.

Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by the manufacturer. PASTORAL AG Strikelab Spray-On Sheep Blowfly Treatment may be applied to all breeds of sheep to obtain protection against fly strike. Significant soiling with dags, mud, dirt or urine may reduce efficacy. Where significant dags are present, a light crutching is indicated prior to treatment. Protection period may be reduced if sheep scour heavily following application.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

WOOL: DO NOT USE less than 3 months before shearing or fibre collection.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 63 days before slaughter for export. Before using this product, confirm the current ESI from the distributor on (03) 5820 8444 or the APVMA website (www.apvma.gov.au/residues).

Use of this product may result in wool residues that do not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

SAFETY DIRECTIONS: Harmful if swallowed. May irritate skin. Will damage eyes. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID INSTRUCTIONS: First aid is not generally required. If in doubt, contact a Poisons Information Centre (e.g. phone Australia 13 11 26).

ADDITIONAL USER SAFETY: Sheep Rehhandling Interval: After treatment wait until sheep are dry before rehhandling. Additional information is listed in the safety data sheet.

ENVIRONMENTAL PROTECTION: DO NOT contaminate dams, rivers, streams or other waterways with the product or used containers.

DISPOSAL: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals onsite. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE: KEEP OUT OF REACH OF CHILDREN. Store below 30°C (room temperature). Protect from frost. Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container.

CONDITIONS OF SALE: The Hunter River Company Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on PASTORAL AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of The Hunter River Company Pty Ltd has any authority to add to or alter these conditions.

B:

EXP:

