

85mm

185mm

#### ENVIRONMENTAL PROTECTION

Doramectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

When used according to the label instructions, PASTORAL AG Doralab Injection is not likely to have an unintended effect that is harmful to the environment or have an overall adverse impact on dung beetle populations. Increased mortality and impaired development of larvae may be expected, but for a limited period after treatment.

#### DISPOSAL

Dispose of container by wrapping with paper and putting in garbage.

#### STORAGE

Store below 30°C (room temperature).

Wash contacted areas thoroughly after handling.  
Do not eat, drink or smoke when using this product.  
**IF SWALLOWED:** Call a POISON CENTRE or doctor if you feel unwell. Rinse mouth. Do NOT induce vomiting.  
Protect from sunlight.  
Store in a dry place. Store in a closed container.

B:

EXP:



MADE IN AUSTRALIA

For the treatment and control of doramectin-sensitive internal and external parasites of cattle and pigs.



**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS  
FOR ANIMAL TREATMENT ONLY

PASTORAL / AG™

**DORALAB™**  
INJECTION  
FOR CATTLE AND PIGS

ACTIVE CONSTITUENT: 10 mg/mL DORAMECTIN

**500 ML**

READ THE CARTON FOR FULL INSTRUCTIONS.

#### DIRECTIONS FOR USE

##### Restraints:

**DO NOT USE** in lactating cows or within 60 days of calving where milk may be used or processed for human consumption.

**Re-treatment interval: DO NOT re-treat animals for 28 days after last treatment.**

##### Dosage and Administration:

**Use contents within 28 days of first broaching of the vial. Store upright. Discard the unused portion. Use a multi-dose syringe for multiple treatments to minimise the number of broachings.**

##### Dose Rate:

CATTLE – 1 mL per 50 kg of bodyweight SC

PIGS – 1 mL per 33 kg (0.3 mL per 10 kg) of bodyweight IM

Administer into the neck region, preferably high up behind the ear. Refer to the dosage tables on the carton as a guide.

#### WITHHOLDING PERIODS

##### MEAT:

CATTLE: 42 days.

PIGS: 35 days.

##### MILK:

**DO NOT USE** in lactating cows or within

60 days of calving.

#### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): 42 days.**

#### SAFETY DIRECTIONS

Poisonous if swallowed. Avoid contact with skin.

#### FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

267931F-3

The Hunter River Company Pty Ltd | ACN 133 798 615 | 74-76 Drummond Road, Shepparton VIC 3630 | Ph: (03) 5820 8444 | APVMA Approval No. 90253/128255