

SPRAY-ON SHEEP BLOWFLY TREATMENT

ACTIVE CONSTITUENT: 65 g/L DICYCLANIL

For the protection of sheep for up to 29 weeks against strike by dicyclanilsusceptible strains of blowfly (Lucilia cuprina). For the protection of mulesing and marking wounds on sheep against strike by dicyclanilsusceptible strains of blowflies during the wound healing process. PAG Strikelab Extra contains 65 g/L of dicyclanil offering premium season long protection against flystrike.

TARGET PARASITES Blowflies (Lucilia cuprina)

DOSAGE & **ADMINISTRATION**


applied as a low-volume protection is required. When . applying PAG Strikelab Extra, the nozzle of the applicator gun should be high enough above the fleece to achieve a 15 cm wide band at each strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool.

WITHHOLDING PERIODS **MEAT:** DO NOT USE less than 14 days before slaughter for human consumption.

FOR THE PREVENTION OF BODY STRIKE (DIAGRAM 1)					
	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS		
10-20	10	10	1		
21-30	12	12	1		
31-50	15		2		
>50	18	9	2		
FOR THE PREVENTION OF BREECH STRIKE (DIAGRAM 1)					
WEIGHT (kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS		
	TOTAL VOLUME/	VOLUME PER	NO. OF		
(kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF		
(kg) 10-20	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF		
(kg) 10-20 21-30	TOTAL VOLUME/ SHEEP (mL) 10 12	VOLUME PER TREATMENT BAND (mL) 10 12	NO. OF		

FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (DIAGRAM 2)					
WEIGHT (kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS		
Up to 30	8	4	2		
31-50	10	5	2		

FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (DIAGRAM 3)					
WEIGHT (kg)	TOTAL VOLUME/ SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS		
Ram lambs	8	4	2		
Ewe lambs	4	4	1		



MILK: DO NOT USE on sheep which are producing or may in the future produce milk that may be used or processed for human

0

WOOL: DO NOT USE less than 3 months before shearing or fibre collection.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

EXPORT SLAUGHTER INTERVAL (ESI)

DO NOT USE less thank 9 weeks before slaughter for export.

RECOMMENDED APPLICATORS

PAG 30ml Drench/Pour-on applicator with spray nozzle.

STORAGE INSTRUCTIONS

Store below 30°C (room temperature). Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.

AVAILABLE SIZES 20 L

For more information, visit pastoralag.com.au For product enquiries, email us at enquiries@pastoralag.com.au or call us on (03) 5820 8444









