

FRONT

100mm




CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

EXACTA
DUAL ACTION ORAL DRENCH FOR SHEEP

Active Constituents:
82.5 g/L LEVAMISOLE HYDROCHLORIDE (equivalent to 70.0 g/L LEVAMISOLE)
36 g/L ALBENDAZOLE OXIDE (equivalent to 34.0 g/L ALBENDAZOLE)

For the control of gastrointestinal roundworms sensitive to benzimidazoles and levamisole including those resistant to either benzimidazoles or levamisole.
For control of lungworm and tapeworm. Reduces the output of viable worm eggs.

10 L

200mm

LEFT SIDE

EFFICACY SPECTRUM (MATURE & IMMATURE)
Including benzimidazole or levamisole resistant worms. This product is not effective against worms that are resistant to BOTH benzimidazole and levamisole compounds.

Roundworms:
Haemonchus contortus – Barber's pole worm
Ostertagia spp. – Small brown stomach worms
Trichostrongylus spp. – Black scour worms
Trichostrongylus axei - Stomach hairworm
Cooperia spp. – Small intestinal worms
Nematodirus spp. – Thin-necked intestinal worms
Oesophagostomum venulosum – Large bowel worm
Charbertia ovina - Large mouth bowel worm
Oesophagostomum columbianum - Nodule worm
Bunostomum spp. - Hookworms

Lungworm:
Dictyocaulus filaria - Large lung worm

Tapeworm:
Moniezia spp. - Heads and segments

Ovicidal:
Roundworm and lungworm eggs


DIRECTIONS FOR USE

Restraints:
DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

Precautions:
Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress.
Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching.
The recommended dose should not be exceeded except under veterinary supervision.
Maintain drench gun carefully to ensure accurate dosing.
Before and after use, clean the drenching gun by flushing out with cold water. DO NOT use hot water.

Dosage and Administration:
SHAKE CONTAINER BEFORE USE.
1 mL EXACTA per 10 kg bodyweight.
Drench sheep orally, using an appropriate drench gun. During drenching, regularly check drench gun to ensure it is delivering the correct dose. A representative sample of the animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs. Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two lines may be appropriate, to avoid excessive overdosing.

BODYWEIGHT (KG)	DOSE (ML)	NO. OF SHEEP TREATED
Up to 20	2.0	5000
21 – 25	2.5	4000
26 – 30	3.0	3333
31 – 35	3.5	2857
36 – 40	4.0	2500
41 – 45	4.5	2222
46 – 50	5.0	2000
51 – 55	5.5	1818
56 – 60	6.0	1666
61 – 65	6.5	1538
66 – 70	7.0	1428
71 – 75	7.5	1333



9 348896 000522

100mm

150mm

RIGHT SIDE

General Directions:
Resistance may develop to any drench.
Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench-resistance test be conducted before any drench is used.

WITHHOLDING PERIODS
MEAT: DO NOT USE less than 10 days before slaughter for human consumption
MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

TRADE ADVICE
EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 10 days before slaughter for export. Before using this product, confirm the correct ESI from The Hunter River Company on (03) 5820 8400 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS
Poisonous if swallowed. Harmful if absorbed by skin contact. May irritate eyes and skin. Avoid contact with eyes and skin. Wear rubber gloves. Wash hands after use.

FIRST AID INSTRUCTIONS
If poisoning occurs, contact a doctor or the Poisons Information Centre. Phone Australia 13 11 26.

ADDITIONAL USER SAFETY
Additional information is listed in the safety data sheet (SDS).

ENVIRONMENTAL PROTECTION
Albendazole and levamisole are extremely toxic to aquatic species. DO NOT contaminate dams, rivers or streams or other water ways with the drench or used containers.

BATCH:
EXP:

DISPOSAL
Triple-rinse container into the medicated drench. Do not dispose of undiluted chemicals onsite. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately at the drumMUSTER collection site. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE
Store below 30°C (room temperature) in original container, tightly closed in a dry place.

May cause an allergic skin reaction.
Suspected of damaging the unborn child.
Do not eat, drink or smoke when using this product.
Wear protective gloves, protective clothing and eye or face protection.
Wash contaminated clothing before reuse.
IF SWALLOWED: Immediately call a POISON CENTRE or doctor. Rinse mouth. Do NOT induce vomiting.
Store locked up in a dry and well-ventilated place.
Store in a closed container.